MTN 035 Product Use End Visit Behavioral Survey Rectal Douche

Table of Contents

| Section: | Title: | Pages: |
|----------|--------------------------|--------|
| A | PSYCHOSOCIAL | |
| В | PRODUCT USE & ADHERENCE | |
| С | PRODUCT ACCEPTABILITY | |
| D | PRODUCT-RELATED PROBLEMS | |
| Е | SEXUAL BEHAVIOR | |
| F | SUBSTANCE USE | |
| G | SOCIAL INFLUENCE | |
| Н | RECOMMENDATIONS | |

PROGRAMMER: Logic checks are in *italics*. Skip patterns are in **CAPITALIZED BOLD**.

SECTION A. PSYCHOSOCIAL

A1. First we would like to ask you about your everyday emotions and experiences. **Since you last completed this survey**, how often have you...

[Response options for 1=Never, 2=Almost never, 3=Sometimes, 4=Fairly often, 5=Very often]

- a. Been upset because of something that happened that you didn't expect?
- b. Felt nervous and "stressed out"?
- c. Found that you could not deal with all the things that you had to do?
- d. Gotten angry because of things that happened that were outside of your control?
- e. Felt that you had so many problems that you could not deal with them?
- f. Felt that you were able to successfully handle the important changes occurring in your life?
- g. Felt able to handle your personal problems?
- h. Felt that things were going your way?
- i. Been able to control hassles in your life?
- j. Felt that you were on top of things?
- k. Been able to control the way to spend your time?

SECTION B. Douche Use & Adherence

The following questions refer to your use of the study provided rectal douches/enemas over the past 4 weeks.

You were asked to insert the **study provided rectal douche/enema** in your rectum <u>at least once a week</u> during the past 4 weeks. However, for different reasons, people might have encountered difficulties using the douche/enema.

B0. Thinking about your experience during these past four weeks, in how many of the weeks did you miss a rectal enema/douche application?

- 1. 0 weeks
- 2. 1 week
- 3. 2 weeks
- 4. 3 weeks
- 5. 4 weeks [SKIP TO B4B]

During this 4-week period you were asked to insert the study douche/enema in your rectum before anal sex or to use it at least once in a seven-day period if you didn't have sex during that time. However, for different reasons, people might have difficulties doing this.

B1. How many times did a partner put a penis in your rectum during the past 4 weeks?

[IF B1 = 0, GO TO B4]

B2. **Of those [B1] times**, how many times did you use the study-provided rectal douche/enema BEFORE receptive anal sex?

```
(If you did not use the douche prior to anal sex, please enter '0'.)
__time(s) [IF 0, SKIP TO B3]
```

- B2a. Typically, how long before **receptive anal intercourse** did you use the study-provided **rectal douche**?
 - 1. Less than 15 minutes
 - 2. Between 15 minutes and 30 minutes.
 - 3. Between 30 minutes and 1 hour
 - 4. Between 1 and 2 hours
 - 5. Between 2 and 3 hours
 - 6. Between 3 and 4 hours
 - 7. 4 hours or more
- B2b. Typically, how many study douches/enemas did you apply before each receptive anal sex occasion?
 - 1. One enema [GO TO B2e]
 - 2. Two enemas
 - 3. Three enemas

- 4. Four enemas
- 5. Five or more enemas

B2c. [If B2b > 1] When you used multiple douches/enemas before receptive anal sex, how often did you rinse the tip with water before reinserting it?

- 1. Never
- 2. Rarely
- 3. Sometimes
- 4. Often
- 5. Always

B2d. [If B2b > 1] When you used multiple douches/enemas before receptive anal sex, how often did you wash the tip with a soap or detergent before reinserting it?

- 1. Never
- 2. Rarely
- 3. Sometimes
- 4. Often
- 5. Always

B2e. What motivated you to use the study provided **rectal douche/enema** BEFORE **receptive anal intercourse**? *Indicate all that apply*

- 1. To follow the study procedures
- 2. I usually use a douche/enema before receptive anal intercourse
- 3. I wanted to be clean
- 4. My sex partner suggested that I use a douche/enema before sex
- 5. Other, please specify: _____

B2f. [**IF B2** ≠ **B1**] Which of the following prevented you from using the study provided rectal douche/enema before sex? (Check all that apply).

- I forgot
- o I did not have the enema bottle with me
- o I didn't have enough time to use the enema before anal sex
- My sexual partner did not want me to use the enema/douche
- o I did not have the privacy needed to use the enema/douche
- I did not like the enema/douche
- The enema/douche made me feel ill
- I was drinking or using drugs
- There was a change in my regular routine
- I felt like people judged me for using a rectal douche
- Other, specify _____

B3. Of those [B1] times that a partner put a penis in your rectum over the past 4 weeks, how many times did you use the study-provided rectal douche AFTER receptive anal sex?

(If you did not use the douche AFTER anal sex, please enter '0'.)

| time(s) [IF 0, SKIP TO B4] |
|---|
| B3a. Typically, how long after receptive anal intercourse did you use a study-provided rectal douche? 1. Less than 15 minutes 2. Between 15 minutes and 30 minutes 3. Between 30 minutes and 1 hour 4. Between 1 and 2 hours 5. Between 2 and 3 hours 6. Between 3 and 4 hours 7. 4 hours or more |
| B3b. Typically, how many douches/enemas did you apply after receptive anal sex? 1. One enema [GO TO B3e] 2. Two enemas 3. Three enemas 4. Four enemas 5. Five or more enemas |
| B3c. [If B3b > 1] When you used multiple douches/enemas after receptive anal sex, how often did you rinse the tip with water before reinserting it? |
| 6. Never7. Rarely8. Sometimes9. Often10. Always |
| B3d. [If B3b > 1] When you used multiple douches/enemas after receptive anal sex, how often did you wash the tip with a soap or detergent before reinserting it? 1. Never 2. Rarely 3. Sometimes 4. Often 5. Always |
| B3e. What made you use a rectal douche/enema AFTER receptive anal intercourse? Indicate all that apply |
| I usually use a douche/enema after receptive anal intercourse I wanted to be clean My sex partner suggested that I use a douche/enema after sex Other, please specify: |
| B4. During the past 4 weeks, how many times did seven days go by in which you did NOT have receptive anal sex? [IF B4 = 0, GO TO B5] |
| B4a. During those [B4] weeks, how many times did you use the study provided rectal douche/enema at least once per week? |

B4b. [**IF B0=5 OR B4a** ≠ **B4**] Which of the following prevented you from using the study provided rectal douche/enema? (Check all that apply).

- I forgot
- I did not have the enema bottle with me
- o My sexual partner did not want me to use the enema/douche
- I did not have the privacy needed to use the enema/douche
- I did not like the enema/douche
- o The enema/douche made me feel ill
- I was drinking or using drugs
- There was a change in my regular routine
- My work/school schedule got in the way
- o I was worried about the long-term effects of the douche on my health
- o I was worried my friends would find out that I was using a rectal douche
- o I was worried my family would find out that I was using a rectal douche
- o I was worried my sexual partners would find out I was using a rectal douche
- o I felt like people judged me for using a rectal douche
- Other, specify _____

B5. Besides the study provided rectal douche/enema, did you use any other rectal products (e.g., douche/enema, suppository) over the past 4 weeks?


- 1. Yes
- 2. No [GO TO B6]

B5a. [If B5=1] Which of the following products did you use? (Check all that apply)

- Over-the-counter disposable enema product (e.g., Fleet®)
- A vaginal douche (e.g., Summer's Eve®) in your rectum
- A non-disposable douche or enema bag system (neoprene or rubber bag, rubber hose, plastic clamp, and plastic or rubber nozzle)____
- A shower head hose and nozzle
- A "sinker", a portable rubber or vinyl hose that attaches to a sink or bidet
- Re-useable bulb enema
- Suppositories (e.g., for hemorrhoids)
- Other (please specify:_____)

B5b. [IF B2 > 0 AND/OR B4a > 0] What percentage of time did you use another rectal douche/enema?

Visual Analog Scale:



| B6. Typically, h | now how far into your rectum did you insert the enema applicator? |
|------------------|---|
| | 1. up to 1 inch |
| | 2. between 1 and 2 inches |
| | 3. between 2 and 3 inches |
| | 4. more than 3 inches |
| B7. Where did | you use a rectal douche? |
| | 1. Toilet |
| | 2. Shower/tub |
| | 3. Sink |
| | 4. Bidet |
| | 5. Bathroom floor |
| | 6. Other, please specify: |
| B8. What temp | perature do you prefer when douching? |
| | 1. Hot |
| | 2. Warm |
| | 3. Cool |
| | 4. No preference |
| B9. In what pos | sition did you prefer to use a rectal douche? |
| | 1. Kneeling |
| | 2. Laying on side |
| | 3. Standing |
| | 4. Squatting or seated over toilet/tub |
| | 5. Other, please specify: |
| | the following did you use to fill the study enema/douche over the past 4 weeks? all that apply) 1. Tap water from the sink |


2. Water provided by study team

| | | Home-made solution (e.g., vinegar) Store-bought product (e.g., vaginal douche, soap) |
|----------------------|---|--|
| 1. 2. 3. 4. | low often did yo Never Rarely Sometimes Often Always | ou need to apply lubricant before inserting the enema/douche bottle? |
| B12. | When you use bowel movem | e a rectal douche, how long does it usually take you before you have a ent? |
| | 1. Occ | eurs immediately |
| | 2. 1-5 | minutes |
| | 3. Mor | e than 5 minutes |
| В13. [| Did you give the 1. Yes 2. No | e study enema/douche to another person to use? [GO TO NEXT SECTION] |
| B14. | How many ene | mas/douches did you give to another person? |
| B15. | 1. A famil 2. A friend 3. A partr 4. A perso 5. A perso | d |
| B16. V | Vhy did you giv | e the study enema/douche to [B15]? |
| SECT | ION C Pactal | Douche ACCEPTABILITY |


3. Water from a public source (e.g., well)

The following questions are about your overall experience with the study douche/enema.


C1. Overall, how much did you like the rectal douche/enema used as part of this study?




C1a. How much did you like how the rectal douche/enema felt inside your rectum immediately after inserting it?




C1b. How much did you like how the rectal douche/enema felt inside your rectum 30 minutes after using it?



C2. If it were available, how likely would you be to continue using the rectal douche/enema provided as part of this study?



C3. How likely would you be to recommend the study provided rectal enema/douche to your friends?



- C4. Compared to other rectal douches/enemas that you may have used in the past, how satisfied were you with the rectal douche/enema used in this study?
 - 1. Very satisfied

| 2. Satisfied | | | | |
|--|------|--|--|--|
| 3. Neutral | | | | |
| 4. Dissatisfied | | | | |
| 5. Very dissatisfied | | | | |
| 6. I had never used a rectal douche/enema in the past | | | | |
| | | | | |
| C5. [IF B2 > 0] Overall , how easy or difficult was it to use the rectal douche BEFORE having receptive anal sex? | 3 | | | |
| 1—2—3—4—5—6—7—8—9—10 | | | | |
| Very Very Easy Difficult | | | | |
| | | | | |
| C6. Overall, how easy or difficult was it to insert the rectal douche/enema? | | | | |
| 1—2—3—4—5—6—7—8—9—10 | | | | |
| Very Easy | | | | |
| Difficult | | | | |
| | | | | |
| C7. The last time you inserted the enema/douche into your rectum, how difficult or easy was to insert? | s it | | | |
| 1—2—3—4—5—6—7—8—9—10 | | | | |
| Very Very Easy Difficult | | | | |
| The following questions are about any changes that you may have experienced while using the study provided rectal douche/enema over the past 4 weeks. | | | | |
| C8. Did your douching practices change over the past 4 weeks? 1. No, I douched as frequently as I typically do 2. Yes, I douched less frequently than I typically do 3. Yes, I douched more frequently than I typically do 4. Not applicable. I had never douched before | | | | |

C9. **Overall**, how did it feel to use the study douche/enema over the past 4 weeks?

1— 2— 3— 4— 5— 6— 7— 8— 9— 10


Very

Very Comfortable
Uncomfortable


C10. Please indicate how hard or easy it would be for you to do each of the following things.

| [1] | [2] | [3] | [4] |
|-----------------|-------------|-------------|-----------------|
| Very Easy to Do | Fairly Easy | Fairly Hard | Very Hard to Do |

- a. To use the rectal douche/enema every time BEFORE having sex?
- b. To discuss your use of the **rectal douche/enema** with a sexual partner BEFORE having sex?
- c. To discuss your use of the rectal douche/enema with a sexual partner AFTER having sex?
- d. To use the rectal douche/enema while under the influence of alcohol or drugs?
- e. To keep the **rectal douche/enema** stored in a private location?
- f. To discretely dispose of the rectal douche/enema after use?
- C11. How much did you like the packaging of the rectal douche/enema?



C12. How easy would it be to carry the rectal douche/enema around if you needed to?



SECTION D. DOUCHE PROBLEMS

Now we would like to ask you about any problems you experienced while using the study enema/douche over the past 4 weeks.

D0. Over the past 28 days, on how many days did you experience any physical discomfort because of the enema/douche?

```
_____ day(s) [IF D0=0, GO TO D1]
```

D0b. Overall, how much did the physical discomfort bother you?

- 1. Not at all
- 2. A little
- 3. Somewhat
- 4. Very much

D1. Did you experience any leakage or soiling after having expelled the water from the douche/enema?

- 1. None [GO TO D2]
- 2. Some
- 3. A lot

D1b. How much were you bothered by the leakage or soiling?

- 1. Not at all
- 2. A little
- 3. Somewhat
- 4. Very much

D2. After having expelled the water from the douche/enema, did you experience any <u>unexpected</u> diarrhea?

- 1. None [GO TO D3]
- 2. Some
- 3. A lot

D2b. How much were you bothered by the unexpected diarrhea?

- 1. Not at all
- 2. A little
- 3. Somewhat
- 4. Very much

D3. After having expelled the water from the douche/enema, did you experience any other <u>unexpected</u> stomach or abdominal problems (such as cramps, bloating, gassiness or passing wind or urge to have a bowel movement)?

- 1. None [GO TO D4]
- 2. Some
- 3. A lot

D3b. How much were you bothered by these stomach or abdominal problems?

- Not at all
- 2. A little
- 3. Somewhat
- 4. Very much

D4. Over the past 28 days, on how many days did you experience any pain because of the enema/douche?

_____ day(s) [IF D4=0, SKIP TO D5]

D4b. Overall, how much did the pain bother you?

- 1. Not at all
- 2. A little
- Somewhat
- 4. Very much

D5. Did the enema/douche cause you any emotional discomfort such as feeling worried, guilty or any other unpleasant feelings?

- 0. No [GO TO D6]
- 1. Yes

D5b. Overall, how much did this emotional discomfort bother you?

- 1. Not at all
- 2. A little
- 3. Somewhat
- 4. Very much

D6. During the four weeks in which you were asked to use the douche/enema, how did it affect your sexual enjoyment?

- 1. My sexual enjoyment was not affected at all
- 2. I had less sexual enjoyment than usual
- 3. I had more sexual enjoyment than usual


D7. [IF B2 > 0] Compared to prior times when you have had receptive anal sex, how different did you feel while having anal sex after using the **study enema/douche** before sex?

- 1. Not at all
- 2. A little
- 3. Somewhat
- 4. Very different

D8. [IF B2 > 0] How did the enema/douche affect your sexual pleasure?

- 1. Increased my sexual pleasure
- 2. Did not change my sexual pleasure
- 3. Decreased my sexual pleasure

D8b. How much were you bothered by having to wait a while to have sex after using the rectal enema/douche?



D9. [IF B2 > 0] How did the douche affect your sex partner's sexual pleasure?

- 1. Increased my partner's sexual pleasure
- 2. Did not change my partner's sexual pleasure
- 3. Decreased my partner's sexual pleasure
- 4. Don't know

D10. Over the past 4 weeks, did any of your partners have a <u>negative</u> reaction regarding your use of the enema/douche?

- 1. No
- 2. Yes, (please specify:_____)

D11. Over the past 4 weeks, did any of your partners have a <u>positive</u> reaction regarding your use of the enema/douche?

- 1. No
- 2. Yes, (please specify:_____)

D12. Over the past 4 weeks, did you have an injury as a result of using the enema/douche?

- 1. Yes [IF NO, SKIP TO NEXT SECTION]
- 2. No

D13. How many times have you had an injury to the anus or rectum while douching in the past 28 days that you used the study product?

- D14. When injuries have occurred, what was the cause? Indicate all that apply
 - 1. Problems with the bottle tip
 - 2. Position played a role
 - 4. Insufficient lubrication
 - 5. Water was too hot
 - 6. Other, please specify: _____

SECTION E. SEXUAL BEHAVIOR

| First, we will ask you questions about your body. |
|--|
| E1. Do you currently have a penis? |
| 1 Yes [SKIP TO E3] |
| 2 No |
| [IF REFUSE TO ANSWSER, ask:] |
| You selected "Refuse to answer." If you do not answer this question, you may be asked questions about your sexual behavior that do not apply to you. Please confirm your answer. |
| 1. Yes, I refuse to answer |
| 2. No, I want to change my answer |
| E2. Do you currently have a vagina? |
| 1 Yes |
| 2 No |
| [IF REFUSE TO ANSWSER, ask:] |
| You selected "Refuse to answer." If you do not answer this question, you may be asked questions about your sexual behavior that do not apply to you. Please confirm your answer. |
| 1. Yes, I refuse to answer |
| 2. No, I want to change my answer |
| RECENT SEXUAL BEHAVIOR |
| E3. In the past 30 days, how many people have you had sex with? (Sex can include anal sex vaginal sex, oral sex, or analingus). |
| [IF REFUSE TO ANSWSER, ask:] |
| You selected "Refuse to answer." If you do not answer this question, you may be asked |

questions about your sexual behavior that do not apply to you. Please confirm your answer.

1. Yes, I refuse to answer

2. No, I want to change my answer

| [If E3=1, ask 4.1. If E3> 1, ask 4.2.] |
|---|
| E4.1. You said you had sex with <i>one</i> person in the past 30 days. Please indicate whether this person was a |
| 1 Man |
| 2 Woman |
| 3 Transgender man |
| 4 Transgender woman |
| [IF REFUSE TO ANSWSER, ask:] |
| You selected "Refuse to answer." If you do not answer this question, you may be asked questions about your sexual behavior that do not apply to you. Please confirm your answer. |
| 1. Yes, I refuse to answer |
| 2. No, I want to change my answer |
| E4.2. You said you had sex with [E4] people in the past 30 days. Please indicate how many of these people were men, women, or transgender. Note that your answers must add up to [Q4]. |
| 1 Men |
| 2 Women |
| 3 Transgender man |
| 4 Transgender woman |
| |
| [IF REFUSE TO ANSWSER, ask:] |
| You selected "Refuse to answer." If you do not answer this question, you may be asked questions about your sexual behavior that do not apply to you. Please confirm your answer. |
| 1. Yes, I refuse to answer |
| |

2. No, I want to change my answer

| [SKIP PATTERNS WILL BE BASED ON ANSWERS TO ANATOMY AND PARTNER'S GENDER] |
|--|
| E5. During the past 30 days, how many times did you have vaginal sex , when <u>you</u> put your penis into a partner's vagina ? [RANGE 0-900] |
| E6. Of the [E5] times you had vaginal sex , how many times did <u>you</u> put your penis into a partner's vagina <u>without a condom</u> , even for a little while? |
| E7. During the past 30 days, how many times did you have insertive anal sex , when <u>you</u> put your penis into a partner's rectum ? |
| E8. Of the [E7] times you had insertive anal sex , how many times did <u>you</u> put your penis into a partner's rectum <u>without a condom</u> , even for a little while? |
| E9 During the past 30 days, how many times did you have receptive anal sex , when <u>a partner's penis was in your rectum?</u> |
| E10. Of the [E9] times you had receptive anal sex , how many times was <u>a partner's penis in your rectum without a condom, even for a little while?</u> |
| Now we would like to ask you about oral sex: |
| E11. During the past 30 days, how many times did <u>you</u> put your mouth on a partner's penis ? |
| E12. During the past 30 days, how many times did <u>a partner</u> put their mouth on your penis ? |
| E13. During the past 30 days, how many times did <u>you</u> put your mouth on a partner's anus ? |
| |

| E14. During the past 30 days, how many times did <u>a partner</u> put their mouth on your anus ? |
|---|
| E15. During the past 30 days, how many times did <u>you</u> put your mouth on a partner's vagina ? |
| E16. During the past 30 days, how many times did <u>a partner</u> put their mouth on your vagina ? |
| E17. During the past 30 days, how many times did you have vaginal sex , when <u>a partner's</u> penis was in <u>your vagina?</u> |
| E18. Of the [E17] times you had vaginal sex , how many times was <u>a partner's penis in your vagina <u>without a condom</u>, even for a little while?</u> |
| The next two questions refer to exchanging sex for money or other goods or services. Remember, your answers are confidential and will not be viewed by clinic staff. |
| E19. During the past 30 days, how many times did you <u>receive</u> money or other goods or services in exchange for sex? |
| E20. During the past 30 days, how many times did you <u>pay</u> money or provide other goods or services in exchange for sex? |
| E21. Please leave us any comments you have about this sexual behavior questionnaire, especially if you thought these questions did not apply to you: |
| |

SECTION F. SUBSTANCE USE

The following questions refer to alcohol and drug use. Remember, your answers are confidential and will not be viewed by clinic staff.

F1. Now you will see a list of different drugs. During the last 30 days, how often have you used each of the following substances?

GO DOWN "A" COLUMN FIRST. IF "0" FOR ANY SUBSTANCES, SKIP B FOR THOSE PARTICULAR SUBSTANCES.

| | [A] | [B] |
|--|---|---|
| | Number of times used in past 30 days | Number of times used before or during sex in past 30 days |
| | | [Only ask if A > 0] |
| a. Tobacco (cigarettes, cigars, chew, snuff, ecigs) | times | times |
| b. Alcohol (beer, wine, liquor) | times | times |
| c. Marijuana/hashish/pot/weed/chamba | times | times |
| d. Synthetic Marijuana/K2/Spice | times | times |
| e. Ecstasy/MDMA | times | times |
| f. Crystal Meth/amphetamines/ methamphetamines/speed/crank/ice/yaba | times | times |
| G. Ketamine/special K | times | times |
| H. GHB (Gamma hydroxybutyrate) | times | times |
| I. Other hallucinogens/LSD/ mushrooms | times | times |
| J. Poppers/amyl nitrate/butyl nitrate | times | times |
| K. Crack | times | times |
| L. Cocaine (not crack) | times | times |
| M. Opium/Opioids (Heroin, Codeine, Vicodin, OxyContin) | times | times |
| N. Mandrax/Quaalude | times | times |
| O. Kratom | times | times |
| P. Khat | times | times |
| Q. Any other pharmaceutical drugs not prescribed to you by a physician | times | times |
| R. Other, please specify: | times | times |

F2. [IF F1A_b>0] Thinking about the times you used alcohol during the <u>last 30 days</u>, how much did you typically use?

- 1. Too little to feel any effect
- 2. Enough to feel it a little
- 3. Enough to feel it a lot
- 4. Enough to get drunk
- 5. Enough to feel like you might pass out
- 6. Enough to pass out

SECTION G. Discussion with Social Network

G1. Have you talked about this study with anyone in your social network?

- 1. No
- 2. Yes (GO TO G1B)

G1a. Different circumstances may prevent people from discussing participation in the study with social networks. Thinking about your experience since you last completed the survey, please choose all the reasons that have made it difficult to talk about the study?

- Embarrassment
- o Fear of Violence
- Potential Misunderstanding
- Don't think it's important
- Lack of Trust
- Stigmatization
- Shyness
- Awkwardness
- Fear of job loss
- o Fear of losing housing
- Fear of criminalization

G1b. Who did you talk about it with? (Check all that apply)

- Friends
- Family
- Sex partners
- Coworkers
- Roommates
- Other _____

G2a. Have you talked about the rectal douche/enema used in this study with anyone in your social network?

- Yes
- No (GO TO NEXT SECTION)

G2b. Who did you talk about it with? (Check all that apply)

- Friends
- Family
- Sex partners
- Coworkers
- Roommates
- Other _____

G2c. How comfortable did you feel talking about the rectal enema/douche?


| {NOTE: LIST | Very | Uncomfortable | Neutral | Comfortable | Very |
|-----------------|---------------|---------------|---------|-------------|-------------|
| FILLS BASED ON | Uncomfortable | | | | Comfortable |
| G2A answers} | | | | | |
| Friends | | | | | |
| Family | | | | | |
| Sex Partner | | | | | |
| Co-Workers | | | | | |
| Roommates | | | | | |
| Fill-in from #2 | | | | | |

G2d. How did they respond when you talked about the rectal enema/douche?

| {NOTE: LIST FILLS | Negatively | Somewhat Negatively | Neutral | Somewhat Positively | Positively | N/A |
|----------------------|------------|------------------------|---------|---------------------|------------|-----|
| BASED ON | | | | | | |
| G2A | | | | | | |
| answers} | | | | | | |
| Friends | | | | | | |
| Family | | | | | | |
| Sex Partner | | | | | | |
| Co-Workers | | | | | | |
| Roommates | | | | | | |
| Fill-in from | | | | | | |
| #2 | | | | | | |

SECTION H. RECOMMENDATIONS

H1. Think about the positive and negative experiences you have had using the rectal douche/enema during the past 4-week period. If this douche/enema was available and it provided some protection against HIV, how likely would you be to use it before receptive anal sex?



| 12. What were the things you liked the most? | |
|--|--|
| • | |
| · | |
| · | |
| | |
| 3. What were the things you liked the <u>least</u> ? | |
| • | |
| | |
| | |

Thank you for completing this questionnaire! Please notify the study staff that you are finished.

END OF QUESTIONNAIRE