Adolescent Contraception

Experiences at Emavundleni Research
Centre
Cape Town



Overview

- Stats
 - o At our site
 - o In SA
- Experiences with adolescent contraception
 - o Barriers
 - o Myths
 - Misconceptions
 - o Adherence
- Clinical Guidance and Legislation in the Western Cape
- Adults vs Adolescents
- Our GOAL
- Discussion & Questions



Statistics

- Use of contraception in married women = 59.8%
 - o Injectable = 28.4%
 - o Sterilization = 15%
 - o Oral contraceptive = 10.9%
 - Male condoms = 4.3%
 - o IUD = less than 1%
 - Implant was not available at the time
- At Emavundleni Research Centre: A quick look at one of our adolescent studies:
 - 15 sexually active young girls
 - 100% of sexually active girls using contraception (protocol requirement)
 - Mostly Injectable: 86%
 - No IUD
 - No pill
 - 2 Implants

Population Reference Bureau, Family Planning Worldwide 2013



Experiences with Adolescent Contraception

- General Barriers (especially with IUD and Implant):
 - Fear of bleeding changes
 - Fear of discovery by parents:
 - The IUD is extremely unpopular among older women in the community we serve.
 - Lack of support by local clinics for IUD users in terms of bleeding changes and management
 - Hesitance of health professionals to offer IUD to adolescent girls
 - Lack of EDUCATION about OPTIONS and CHOICE
 - Need among young women to prove their fertility (amenorrhea is a problem)



INJECTABLE

- DMPA and Net-En available at our site
- Myths and Misconceptions:
 - Injectable is the only choice. Either 3 monthly or 2 monthly. Most girls choose 3 monthly because it meant less visits to the clinic
 - Amenorrhea: Girls who develop amenorrhea stop the method, believing that their blood is "blocked" or "building up". They can eventually smell the blood and it comes out as nosebleeds.

Adherence

- Nurses at the clinic are unfriendly. They get reprimanded if they are late for their injection so they just don't go back.
- Clinic times vs school times



Implant

- New Urban Myth:
 - Gangsters are attacking women and cutting the implant out in order to smoke it= massive barrier
 - Women have requested to remove implants soon after insertion because of this
 - Not confirmed yet
- Fear of insertion process
- Implant damages your arm
- Belief that long term methods are only for older women



IUD

Myths and Misconceptions

- The IUD is only indicated for women who have completed their families
- Hesitation by health professionals to offer IUD to 16-18 year olds
- o IUD causes STIs
- o It moves around the lower abdomen

Barriers

- Mothers of young girls force them to remove the IUD
- No support from local clinics
- Fear of pain and prolonged bleeding
- Beliefs and attitudes of health care workers



Clinical Guidance

- No specific guidance for adolescent contraception
- Our site has a general contraception SOP but nothing specific to adolescents
- No laws governing adolescent contraception provision
- WHO Medical Eligibility Criteria: no medical reason currently exists for denying contraception based on age



Adults vs Adolescents

- Factors that influence adolescents: school, friends parents
- Counseling messages need to be clear
- Dispel myths
- Provide support for side effects specifically bleeding changes
- Education for both adolescents and adults to make informed choices
- We would like to increase implant and IUD uptake



Thank You

